

Guoging

Achievements

The best graduate researcher award from Department of Physics is Mr Chang Guoqing. Mr Chang's research focuses on varies topological phases in condensed matter physics, such as topological semimetals and topological insulators. He made important contributions in several projects related to the topological materials and prepared numbers of papers where Mr Chang is first author. Mr Chang's outstanding research has been published in top journals such as Science, Science Advances, PNAS, Nature Materials, Nature Physics, Nature Communications, and Physical Review Letters. With an essential blend with his deep theoretical understanding and computation skills, Mr Chang is very helpful for experimentalists and plays the key role in the discovery of several new topological materials.

Awards

- ✓ Best Graduate Researcher Award 2017
- Chinese Government Award for Outstanding Self-financed PhD Student Abroad (2016)



Research Areas

I am working on varies topological phases in condensed matter physics, such as topological semimetals and topological insulators.

I have predicted several Weyl semimetal candidates: TaAs class, LaAlGe family, W1-xMoxTe2, Ta3S2, Co2TiX and SrSi2. Among them, TaAs family, LaAlGe, and W1-xMoxTe2 have been experimentally confirmed.

Besides searching for new materials, I am also working on classifying previously uncharacterized new topology phases. Specifically, in chiral space groups, I proposed the theory of Kramers Weyl fermions. I have also defined new Fermions beyond the Standard Model, such as three-fold Nexus fermions and Hopf link semimetals.

My research involves extensive and close collaborations with distinguished many experimental groups. I am experienced in analyzing data from angle-resolved photoemission (ARPES), and scanning tunneling spectroscopy (STM). My theoretical calculations of bulk electronic structure. surface states, interference patterns, and so on are always offering crucial and timely support for experimental measurements of our collaborators.

Discovery of a Weyl fermion semimetal and topological Fermi arcs

RESEARCH | RESEARCH ARTICLES

NIH (8UL1TR000043); the French National Research Agency (ANR) under the "Investments for the Future" program (gran ANR-10-IAHU-01), grant IFNGPHOX (13-ISV3-0001-01 to J.B.), and grant GENCMCD (11-BSV3-005-01 to A.P); Laboratoire d'Excellence Integrative Biology of Emerging Infectious Diseases (ANR-10-LABX-62-IBEID); the National Health and Medical Research Council (NHMRC) (to E.K.D., C.S.M., S.G.T, and J.Mc.) the St. Giles Foundation; the National Institute of Allergy and Infectious Diseases (R37AI095983 to J.-L.C.); and the NIH

(contract HHSN272200900044C to A.S.). S.O. was supported to

the Promotion of Science (25713039 and 25670477), J.G.M. by the Canadian Institutes of Health Research, R.M.-B. by the European Molecular Biology Organization, Y.I. by the AXA Research Fund, Development Award, and F.S. by grants from the European Research Council (323183 PREDICT) and the Swiss National Biomedicine and the Center of Medical Immunology are supported by the Helmut Horten Foundation. S.A.-M. is the bronchial asthma research chair of the Prince Naif Center for Immunology Research.

SUPPLEMENTARY MATERIALS Materials and Methods Supplementary Text Figs. S1 to S19 Tables S1 to S3 References (38-67) 7 December 2014; accepted 29 June 2015 Published online 9 July 2015

TOPOLOGICAL MATTER

Discovery of a Weyl fermion semimetal and topological Fermi arcs

Su-Yang Xu,^{1,2*} Ilya Belopolski,^{1*} Nasser Alidoust,^{1,2*} Madhab Neupane,^{1,3*} Guang Bian, Chenglong Zhang, Raman Sankar, Guoqing Chang, 7 Zhujun Yuan, Chi-Cheng Lee, ^{6,7} Shin-Ming Huang, ^{6,7} Hao Zheng, ¹ Jie Ma, ⁸ Daniel S. Sanchez, ¹ BaoKai Wang, ^{6,7,9} Arun Bansil, ⁹ Fangcheng Chou, ⁵ Pavel P. Shibayev, ^{1,10} Hsin Lin, ^{6,7} Shuang Jia, 4,11 M. Zahid Hasan 1,2 †

A Weyl semimetal is a new state of matter that hosts Weyl fermions as emergent quasiparticles and admits a topological classification that protects Fermi arc surface states on the boundary of a bulk sample. This unusual electronic structure has deep analogies with particle physics and leads to unique topological properties. We report the experimental discovery of a Weyl semimetal, tantalum arsenide (TaAs). Using photoemission spectroscopy, we directly observe Fermi arcs on the surface, as well as the Weyl fermion cones and Weyl nodes in the bulk of TaAs single crystals. We find that Fermi arcs terminate on the Weyl fermion nodes, consistent with their topological character. Our work opens the field for the experimental study of Weyl fermions in physics and materials science.

I fermions have long been known in | to points of bulk band degeneracy, Weyl nodes,

quantum field theory, but have not been which are associated with a chiral charge that

nature (1-3). Recently, it was understood that a Weyl fermion can emerge as a quasiparticle in certain crystals, Weyl fermion semimetals (1-22). Despite being a gapless metal, a Wevl semimetal is characterized by topological invariants, broadening the classification of topological phases of matter beyond insulators. Specifically, Weyl fermions at zero energy correspond ¹Laboratory for Topological Quantum Matter and Spectroscopy (B7), Department of Physics, Princeton University, Princeton, NJ 08544, USA. ²Princeton Center for Complex Materials, Princeton Institute for Science and Technology of Materials, Princeton University, Princeton, NJ

observed as a fundamental particle in

08544, USA. 3Condensed Matter and Magnet Science Group Los Alamos National Laboratory, Los Alamos, NM 87545, USA. ⁴International Center for Quantum Materials, School of Physics, Peking University, China. ⁵Center for Condensed Matter Sciences, National Taiwan University, Taipei 10617, Taiwan. ⁶Centre for Advanced 2D Materials and Graphene Research Centre National University of Singapore, 6 Science Drive 2. Singapore 117546. Department of Physics, National University of Singapore, 2 Science Drive 3, Singapore 117542. 8 Quantum Condensed Matter Division, Oak Ridge National Laboratory, Oak Ridge, TN 37831, USA. ⁹Department of Physics, Northeastern University, Boston, MA 02115, USA ¹⁰Princeton Institute for Science and Technology of Materials, Princeton University, Princeton, NJ 08544, USA. ¹Collaborative Innovation Center of Quantum Matter, Beijing 100871, China. These authors contributed equally to this work. †Corresponding protects gapless surface states on the boundary of a bulk sample. These surface states take the form of Fermi arcs connecting the projection of bulk Weyl nodes in the surface Brillouin zone (BZ) (6). A band structure like the Fermi arc surface states would violate basic band theory in an isolated two-dimensional (2D) system and can only arise on the boundary of a 3D sample, providing a dramatic example of the bulk-boundary correspondence in a topological phase. In contrast to topological insulators where only the surface states are interesting (21, 22), a Weyl semimetal features unusual band structure in the bulk and on the surface. The Weyl fermions in the bulk are predicted to provide a condensedmatter realization of the chiral anomaly, giving rise to a negative magnetoresistance under parallel electric and magnetic fields, unusual optical conductivity, nonlocal transport, and local nonconservation of ordinary current (5, 12-16). At the same time, the Fermi arc surface states are predicted to show unconventional quantum oscillations in magneto-transport, as well as unusual quantum interference effects in tunneling spectroscopy (17-19). The prospect of the realization of these phenomena has inspired much experimental and theoretical work (1-22). Here we report the experimental realization of a Weyl semimetal in a single crystalline material,

SCIENCE sciencemag.org

author, E-mail: mzhasan@princeton.edu

and soft x-ray (SX) angle-resolved photoemission spectroscopy (ARPES), we systematically and differentially study the surface and bulk electronic structure of TaAs. Our ultraviolet (low-photonenergy) ARPES measurements, which are highly surface sensitive demonstrate the existence of the Fermi arc surface states, consistent with our band calculations presented here. Moreover, our SX-ARPES measurements, which are reasonably bulk sensitive, reveal the 3D linearly dispersive bulk Weyl cones and Weyl nodes. Furthermore, by combining the low-photon-energy and SX-ARPES data, we show that the locations of the projected bulk Weyl nodes correspond to the terminations of the Fermi arcs within our experimental resolution. These systematic measurements demonstrate TaAs as a Weyl semimetal. The material system and theoretical considerations

tantalum arsenide (TaAs). Using the combina-

tion of the vacuum ultraviolet (low-photon-energy)

Tantalum arsenide is a semimetallic material that crystallizes in a body-centered tetragonal lattice system (Fig. 1A) (23). The lattice constants are a = 3.437 Å and c = 11.656 Å, and the space group is $I4_1md$ (#109, C_{4v}), as consistently reported in previous structural studies (23-25). The crystal consists of interpenetrating Ta and As sublattices, where the two sublattices are shifted by $(\frac{a}{2}, \frac{a}{2}, \delta)$, $\delta \approx \frac{c}{12}$. Our diffraction data match well with the lattice parameters and the space group $I4_1md$ (26). The scanning tunneling microscopic (STM) topography (Fig. 1B) clearly resolves the (001) square lattice without any obvious defect. From the topography, we obtain a lattice constant a = 3.45 Å. Electrical transport measurements on TaAs confirmed its semimetallic transport properties and reported negative magnetoresistance, suggesting the anomalies due to

Weyl fermions (23). We discuss the essential aspects of the theoretically calculated bulk band structure (9, 10) that predicts TaAs as a Weyl semimetal candidate. Without spin-orbit coupling, calculations (9, 10) show that the conduction and valence bands interpenetrate (dip into) each other to form four 1D line nodes (closed loops) located on the k_x and k_y planes (shaded blue in Fig. 1, C and E). Upon the inclusion of spin-orbit coupling, each line node loop is gapped out and shrinks into six Wevl nodes that are away from the $k_v = 0$ and $k_u = 0$ mirror planes (Fig. 1E, small filled circles). In our calculation, in total there are 24 bulk Weyl cones (9, 10), all of which are linearly dispersive and are associated

RESEARCH | RESEARCH ARTICLES

because it arises from the nonzero chiral charge (Fig. 3A). We now choose an incident photon energy (i.e., a k_z value) that corresponds to the k-space of the projected bulk Weyl nodes (26), which in this case is ± 2 . Therefore, this property distinguishes the crescent Fermi arcs not only from any closed contour but also from the double Fermi arcs in Dirac semimetals (27, 28), because the bulk Dirac nodes do not carry any net chiral charges (26). After observing the surface electronic structure containing Fermi arcs in our ARPES data, we are able to slightly tune the free parameters of our surface calculation and obtain a calculated surface Fermi surface that reproduces and explains our ARPES data (Fig. 1G). This serves as an important cross-check that our data and interpretation are self-consistent. Specifically, our surface calculation indeed also reveals the crescent Fermi arcs that connect the projected W2 Weyl nodes near the midpoints of each $\overline{\Gamma}$ – \overline{X} or $\overline{\Gamma}$ – \overline{Y} line (Fig. 1G). In addition, our calculation shows the bowtie surface states centered at the \overline{X} point, also consistent with our ARPES data. According to our calculation, these bowtie surface states are in fact Fermi arcs (26) associated with the W1 Wevl nodes near the BZ boundaries. However, our ARPES data cannot resolve the arc character because the W1 Weyl nodes are too close to each other in momentum space compared to the experimental resolution. Additionally, we note that the agreement between the ARPES data and the surface calculation upon the contour at the \overline{Y} point can be further improved by fine-optimizing the surface parameters. To establish the topology, it is not necessary for the data to have a perfect correspondence with the details of calculation because some changes in the choice of the surface potential allowed by the free parameters do not change the topology of the materials, as is the case in topological insulators (21, 22). In principle, Fermi arcs can coexist with additional closed contours in a Weyl semimetal (6, 9), just as Dirac cones can coexist with additional trivial surface states in a topological insulator (21, 22). Partic-

Bulk measurements

We now present bulk-sensitive SX-ARPES (29) data, which reveal the existence of bulk Weyl cones and Weyl nodes. This serves as an independent proof of the Wevl semimetal state in TaAs. Figure 3B shows the SX-ARPES measured $k_x - k_z$ Fermi surface at $k_y = 0$ (note that none of the Weyl nodes are located on the $k_y = 0$ plane). We emphasize that the clear dispersion along the k_r direction (Fig. 3B) firmly shows that our SX-ARPES predominantly images the bulk bands. SX-ARPES boosts the bulk-surface contrast in favor of the bulk band structure, which can be further tested by measuring the band dispersion along the k_z axis in the SX-ARPES setting. This is confirmed by the agreement between the ARPES data (Fig. 3B) and the corresponding bulk band calculation

ularly, establishing one set of Weyl Fermi arcs

is sufficient to prove a Weyl semimetal (6). This is achieved by observing the crescent Fermi

arcs as we show here by our ARPES data in Fig.

2, which is further consistent with our surface

location of W2 Weyl nodes and map the corresponding $k_r - k_u$ Fermi surface. As shown in Fig. 3C, the Fermi points that are located away from the k_x or k_y axes are the W2 Weyl nodes. In Fig. 3D, we clearly observe two linearly dispersive cones that correspond to the two nearby W2 Wevl nodes along cut 1. The k-space separation between the two W2 Weyl nodes is measured to be 0.08 Å⁻¹, which is consistent with both the bulk calculation and the separation of the two terminations of the

crescent Fermi arcs measured in Fig. 2. The linear dispersion along the out-of-plane direction for the W2 Weyl nodes is shown by our data in Fig. 3E. Additionally, we also observe the W1 Weyl cones in Fig. 3, G to I. Notably, our data show that the ener gy of the bulk W1 Weyl nodes is lower than that of the bulk W2 Weyl nodes, which agrees well with our calculation shown in Fig. 3J and an independent modeling of the bulk transport data on TaAs (23). In general, in a spin-orbit-coupled bulk crystal, point-like linear band crossings can either be Weyl cones or Dirac cones. Because the observed bulk




Fig. 2. Observation of topological Fermi arc surface states on the (001) surface of TaAs. (A) ARPES Fermi surface map and constant-binding energy contours measured with incident photon energy of 90 eV. (B) High-resolution ARPES Fermi surface map of the crescent Fermi arcs. The k-space range of this map is defined by the blue box in (A). (C and D) Energy dispersion maps along cuts I and II. (E) Same Fermi surface map as in (B). The dotted lines define the k-space direction for cuts I and II. The numbers 1 to 6 denote the Fermi crossings that are located on cuts I and II. The white arrows show the evolution of the constant-energy contours as one varies the binding energy, which is obtained from the dispersion maps in (C) and (D). (F) A schematic showing the evolution of the Fermi arcs as a function of energy, which clearly distinguishes between two Fermi arcs and a closed contour. (G) The difference between the constant-energy contours at the binding energy $E_B = 20$ meV and the binding energy $E_B = 0$ meV, from which one can visualize the evolution of the constant-energy contours through $k_v - k_v$ space. The range of this map is shown by the white dotted box in (A).

SCIENCE sciencemag.org

7 AUGUST 2015 • VOL 349 ISSUE 6248 615

RESEARCH

cones in Fig. 3. C and D, are located neither at Kramers' points nor on a rotational axis, they cannot be identified as bulk Dirac cones and have to be Wevl cones according to topological theories (6, 28). Therefore, our SX-ARPES data alone prove the existence of bulk Weyl nodes. The agreement between the SX-ARPES data and our bulk calculation, which only requires the crystal structure and the lattice constants as inputs, provides a further

Bulk-surface correspondence

surface Fermi arc terminations match with the projection of the bulk Weyl nodes on the surface BZ. We superimpose the SX-ARPES measured bulk Fermi surface containing W2 Weyl nodes (Fig. 3C) onto the low-photon-energy ARPES Fermi surface containing the surface Fermi arcs (Fig. 2A) to scale. From Fig. 4A we see that all the arc terminations and projected Weyl nodes match with each other within the k-space region that is covered in our measurements. To establish this point quantitatively, in Fig. 4C, we show the zoomed-in map near the crescent Fermi arc terminations, from which we obtain the k-space location of the terminations to be at $\vec{k}_{arg} = (0.04 \pm 0.01 \,\text{Å}^{-1}, 0.51 \pm 0.01 \,\text{Å}^{-1})$. Figure 4D shows the zoomed-in map of two nearby W2 Weyl nodes, from which we obtain the k-space location of the W2 Weyl nodes to be at \vec{k}_{W2} = $(0.04 \pm 0.015 \,\text{Å}^{-1} \, 0.53 \pm 0.015 \,\text{Å}^{-1})$ In our bulk calculation, the k-space location of the W2 Weyl nodes is found to be at $(0.035\,\text{Å}^{-1}, 0.518\,\text{Å}^{-1})$ Because the SX-ARPES bulk data and the lowphoton-energy ARPES surface data are completely independent measurements obtained with two different beamlines, the fact that they match well provides another piece of evidence of the topological nature (the surface-bulk correspondence) of the Weyl semimetal state in TaAs. In figs. S5 to S7, we further show that the bulk Wevl cones can also be observed in our low-photon-energy ARPES data, although their spectral weight is much lower than the surface state intensities that dominate the data. Our demonstration of the Weyl fermion semimetal state in and Fermi arc surface metals paves the way (30) for the realization of many fascinating topological quantum phenomena.

REFERENCES AND NOTES H. Weyl, Z. Phys. 56, 330-352 (1929). L. Balents, *Physics* 4, 36 (2011).
F. Wilczek, *Phys. Today* 51, 11 (1998). Fundamentals and Perspectives, F. Ortmann, S. Roche,

7. B. Singh et al., Phys. Rev. B 86, 115208 (2012). 8. S.-Y. Xu et al., Science 332, 560-564 (2011). S.-M. Huang et al., Nat. Commun. 6, 7373 (2015).
H. Weng, C. Fang, Z. Fang, F. B. Andrei Bernevig, X. Dai, Phys.

14. C.-X. Liu, P. Ye, X.-L. Qi, Phys. Rev. B 87, 235306 (2013)

SCIENCE sciencemag.org

15. H. Wei, S.-P. Chao, V. Aji, Phys. Rev. Lett. 109, 196403 (2012

16. P. E. C. Ashby, J. P. Carbotte, Phys. Rev. B 87, 245131 (2013).

4. M. Z. Hasan, S.-Y. Xu, M. Neupane, in Topological Insulators H. B. Nielsen, M. Ninomiya, *Phys. Lett. B* 130, 389–396 (1983).
X. Wan, A. M. Turner, A. Vishwanath, S. Y. Savrasov, *Phys. Rev.*

11. J. Liu. D. Vanderbilt, Phys. Rev. B 90, 155316 (2014). 12. A. A. Zyuzin, A. A. Burkov, Phys. Rev. B 86, 115133 (2012). 13. D. T. Son, B. Z. Spivak, Phys. Rev. B 88, 104412 (2013).

 T. Ojanen, Phys. Rev. B 87, 245112 (2013). 20. S. Murakami, New J. Phys. 9, 356 (2007). M. Z. Hasan, C. L. Kane, Rev. Mod. Phys. 82, 3045-3067 . Qi, S.-C. Zhang, Rev. Mod. Phys. 83, 1057-1110 (2011). . C.-L. Zhang et al., Observation of the Adler-Bell-Jackiw chiral anomaly in a Weyl semimetal. http://arxiv.org/abs/1503.

5. S. Furuseth et al., Acta Chem. Scand. 19, 95-106 (1965). 26. Materials and methods are available as supplementary materials on Science Online 28. B.-J. Yang, N. Nagaosa, Nat. Commun. 5, 4898 (2014) V. N. Strocov et al., J. Synchrotron Radiat. 21, 32–44 (2014).
M. Z. Hasan, "Topological Insulators: A New Form of Quantum Matter," presentation at 2014 Nobel Symposium on New Forms of Matter, www.fysik.su.se/~ardonne/nobel/ns156-program.

24. J. J. Murray et al., J. Less Common Met. 46, 311-320 (1976)

ACKNOWLEDGMENTS

Work at Princeton University and Princeton-led synchrotron-base ARPES measurements were supported by the Gordon and Betty Moore Foundations EPiQS Initiative through grant GBMF4547 (Hasan). First-principles band structure calculations at National University of Singapore were supported by the National Research Foundation, Prime Minister's Office, Singapore, under its NRF fellowship (NRF Award no. NRF-NRFF2013-03). Single-crystal

growth was supported by National Basic Research Program of China (grant nos. 2013CB921901 and 2014CB239302) and characterization by U.S. Department of Energy DE-FG-02-05ER46200. F.C. acknowledges the support provided by MOST-Taiwan under project no. 102-2119-M-002-004. We gratefully acknowledge J. D. Denlinger, S. K. Mo, A. V. Fedorov M. Hashimoto, M. Hoesch, T. Kim, and V. N. Strocov for their peamline assistance at the Advanced Light Source, the Stanford Synchrotron Radiation Lightsource, the Diamond Light Source and the Swiss Light Source under their external user programs. art of the work was carried out at the Swiss Light Source throug the external/international facility user program. We thank -R. Chang for help on theoretical band structure calculations. W also thank L. Balents, D. Huse, I. Klebanov, T. Neupert, A. Polyako ². Steinhardt, H. Verlinde, and A. Vishwanath for discussions University, M.Z.H. acknowledges hospitality of the Lawrence a visiting scientist. A patent application is being prepared on behal

SUPPLEMENTARY MATERIALS www.sciencemag.org/content/349/6248/613/suppl/DC1 Supplementary Text References (31, 32) 15 February 2015; accepted 6 July 2015 Published online 16 July 2015

10.1126/science.aaa9297

of the authors on the discovery of a Weyl semimetal.

REPORTS

Rock physics of fibrous rocks akin to Roman concrete explains uplifts at Campi Flegrei Caldera

Tiziana Vanorio* and Waruntorn Kanitpanyacharoen†

Uplifts in the Campi Flegrei caldera reach values unsurpassed anywhere in the world (~2 meters). Despite the marked deformation, the release of strain appears delayed. The rock physics analysis of well cores highlights the presence of two horizons, above and below the seismogenic area, underlying a coupled process. The basement is a calc-silicate rock housing hydrothermal decarbonation reactions, which provide lime-rich fluids. The caprock above the seismogenic area has a pozzolanic composition and a fibril-rich matrix that results from lime-pozzolanic reactions. These findings provide evidence for a natural process reflecting that characterizing the cementitious pastes in modern and Roman concrete. The formation of fibrous minerals by intertwining filaments confers shear and tensile strength to the caprock, contributing to its ductility and increased resistance

he caldera of Campi Flegrei (fig. S1) is one | nearly 40,000 people were evacuated for almost of the active hydrothermal systems of the Mediterranean region experiencing notaarea. During the last crisis of 1982 to 1984,

Stanford Rock Physics Laboratory, Department of Geophysics, Stanford University, Stanford, CA 94305, USA. *Corresponding author. E-mail: tvanorio@stanford.edu †Present address: Department of Geology, Faculty of Science, Chulalongkorn

Puteoli, owing to the large uplifts (~2 m over ble unrest episodes in a densely populated 2 years) and the persistent seismic activity (1). The evacuation severely hampered the economy and the social make-up of the community, which included the relocation of schools and commercial shops, as well as the harbor being rendered useless for docking. Despite the large uplifts, the release of strain appears delayed (2, 3). Seismicity reaches a magnitude of 4.0 only upon relatively

2 years from the main town of Pozzuoli, the Roman

RESEARCH | RESEARCH ARTICLES

with a single chiral charge of ±1 (Fig. 1E). We denote | is implemented as a screw axis that sends the | ize the evolution of the constant-energy con the 8 Weyl nodes that are located on the brown plane $(k_z = \frac{2\pi}{c})$ as W1 and the other 16 nodes that are away from this plane as W2. At the (001) surface BZ (Fig. 1F), the eight W1 Weyl nodes are projected in the vicinity of the surface BZ edges, \overline{X} and \overline{Y} . More interestingly, pairs of W2 Weyl nodes with the same chiral charge are projected onto the same point on the surface BZ. Therefore, in total there are eight projected W2 Weyl nodes with a projected chiral charge of ±2, which are located near the midpoints of the $\overline{\Gamma}$ – \overline{X} and the $\overline{\Gamma}$ – \overline{Y} lines. Because the ± 2 chiral charge is a projected value, the Weyl cone is still linear (9). The number of Fermi arcs terminating on a projected Wevl node must equal its projected chiral charge. Therefore, in TaAs, two Fermi arc surface states must terminate on each

projected W2 Weyl node.

Surface electronic structure of TaAs We carried out low-photon-energy ARPES measurements to explore the surface electronic structure of TaAs. Figure 1H presents an overview of the (001) Fermi surface map. We observe three types of dominant features, namely a crescent-shaped feature in the vicinity of the midpoint of each $\overline{\Gamma} - \overline{X}$ or $\overline{\Gamma} - \overline{Y}$ line, a bowtielike feature centered at the \overline{X} point, and an extended feature centered at the \overline{Y} point. We find that the Fermi surface and the constantenergy contours at shallow binding energies (Fig. 2A) violate the C_4 symmetry, considering the features at \overline{X} and \overline{Y} points. In the crystal structure of TaAs, where the rotational symmetry

crescent-shaped features. Their peculiar shape suggests the existence of two arcs, and their termination points in k-space seem to coincide with the surface projection of the W2 Weyl nodes. Because the crescent feature consists of two nonclosed curves, it can either arise from two Fermi arcs or a closed contour; however, the decisive property that clearly distinguishes one case from the other is the way in which the constant-energy contour evolves as a function of energy. As shown in Fig. 2F, in order for the crescent feature to be Fermi arcs, the two nonclosed curves have to move (disperse) in the same direction as one varies the energy (26). We now provide ARPES data to show that the crescent features in TaAs indeed exhibit this "copropagating" property. To do so, we single out a crescent feature, as shown in Fig. 2, B and E, and show the band dispersions at representative momentum space cuts, cut I and cut II, as defined in Fig. 2E. The corresponding E-kdispersions are shown in Fig. 2, C and D. The evolution (dispersive "movement") of the bands as a function of binding energy can be clearly read from the slope of the bands in the dispersion maps and is indicated in Fig. 2E by the white arrows. It can be seen that the evolution of the two nonclosed curves is consistent with the copropagating property. To further visual-

crystal back into itself after a C_4 rotation and a

translation by $\frac{c}{2}$ along the rotation axis, such an

asymmetry is expected in calculation. The crys-

tallinity of the (001) surface in fact breaks the

rotational symmetry. We now focus on the

namely $\Delta I(k_x, k_y) = I(E_{\rm B} = 20 \text{ meV}, k_x, k_y)$ $I(E_{\rm B}=0~{\rm meV},k_x,k_y)$, where I is the ARPES intensity. The k-space regions in Fig. 2G that to the constant-energy contour at $E_{\rm B}=0\,{\rm meV}$ whereas those regions with positive spectral weight (blue) correspond to the contour at $E_{\rm B}=20$ meV. Thus, one can visualize the two contours in a single k_x , k_y map. The alternating "red-blue-red-blue" sequence for each crescent feature in Fig. 2G shows the copropagating property there are two crescent features, one located near the $k_r = 0$ axis and the other near the $k_r = 0$ axis, in Fig. 2G. The fact that we observe the copropagating property for two independent crescent features that are 90° rotated with respect to each other further shows that this observation is not due to artifacts, such as a k misalignment while performing the subtraction. The above systematic data reveal the existence of Fermi arcs on the (001) surface of TaAs. Just as one can identify a crystal as a topological insulator by observing an odd


tour throughout k_x, k_y space, we use surface

state constant-energy contours at two slightly

different binding energies, namely $E_{\rm B}=0=E_{\rm F}$

and $E_{\rm B}=20\,$ meV. Figure 2G shows the difference

number of Dirac cone surface states, we emphasize that our data here are sufficient to identify TaAs as a Wevl semimetal because of bulkboundary correspondence in topology. Theoretically, the copropagating property of the Fermi arcs is unique to Weyl semimetals



614 7 AUGUST 2015 • VOL 349 ISSUE 624

sciencemag.org SCIENCE

RESEARCH | RESEARCH ARTICLES





Fig. 3. Observation of bulk Weyl fermion cones and Weyl nodes in **TaAs.** (A and B) First-principles calculated and ARPES-measured $k_z - k_x$ Fermi surface maps at $k_y = 0$, respectively. (**C**) ARPES-measured and firstprinciples calculated $k_x - k_y$ Fermi surface maps at the k_z value that corresponds to the W2 Weyl nodes. The dotted line defines the k-space cut direction for cut 1, which goes through two nearby W2 Weyl nodes along the k_{ν} direction. The black cross defines cut 2, which means that the k_{ν}, k_{ν} values are fixed at the location of a W2 Weyl node and one varies the k2 value. (**D**) ARPES $E - k_y$ dispersion map along the cut 1 direction, which clearly shows the two linearly dispersive W2 Weyl cones. (**E**) ARPES $E - k_z$

dispersion map along the cut 2 direction, showing that the W2 Weyl cone also disperses linearly along the out-of-plane k_z direction. (**F**) First principles calculated $E - k_z$ dispersion that corresponds to the cut 2 shown in (E). (G) ARPES measured $k_x - k_y$ Fermi surface maps at the k_z value that corresponds to the W1 Weyl nodes. The dotted line defines the k-space cut direction for cut 3, which goes through the W1 Weyl nodes along the k_y direction. (**H** and **I**) ARPES $E - k_y$ dispersion map and its zoomed-in version along the cut 3 direction, revealing the linearly dispersive W1 Weyl cone. (J) First-principles calculation shows a 14-meV energy difference between the W1 and W2 Weyl nodes.

logically nontrivial state in TaAs. (A) Low-photonenergy ARPES Fermi surface map (hv = 90 eV) from Fig. 2A, with the SX-ARPES map (hv = 650 eV) from Fig. 3C overlaid on top of it to scale, showing that the locations of the projected bulk Wevl nodes correspond to the terminations of the surface Fermi arcs. (B) The bottom shows a rectangular tube in the bulk BZ that encloses four W2 Weyl nodes. These four W2 Weyl nodes project onto two points at the (001) surface BZ with projected chiral charges of ±2, shown by the brown circles. The top surface shows the ARPES measured crescent surface Fermi arcs that connect these two projected Weyl nodes. (C) Surface state Fermi surface map at the k-space region corresponding to the terminations of the crescent Fermi arcs. The k-space region is defined by the black dotted box in (E). (D) Bulk Fermi surface map at the k-space region corresponding to the W2 Weyl nodes. The k-space region is defined by the black dotted box in Fig. 3C. (E) ARPES and schematic of the crescent-shaped co-

Fig. 4. Surface-bulk correspondence and the topo-



propagating Fermi arcs. (F) High resolution ARPES maps of the Fermi arcs and the Weyl fermion nodes.

616 7 AUGUST 2015 • VOL 349 ISSUE 6248

sciencemag.org SCIENCE

Sample-Size Antibod

EXTENDED PDF FORMA

Science MAAAS

Discovery of a Weyl fermion semimetal and topological Fermi arcs Su-Yang Xu et al. Science 349, 613 (2015); DOI: 10.1126/science.aaa9297

This copy is for your personal, non-commercial use only

If you wish to distribute this article to others, you can order high-quality copies for your colleagues, clients, or customers by clicking here. Permission to republish or repurpose articles or portions of articles can be obtained by

following the guidelines here.

The following resources related to this article are available online at www.sciencemag.org (this information is current as of March 13, 2016):

Updated information and services, including high-resolution figures, can be found in the online version of this article at: /content/349/6248/613.full.html

Supporting Online Material can be found at: /content/suppl/2015/07/15/science.aaa9297.DC1.html

A list of selected additional articles on the Science Web sites related to this article can be found at: /content/349/6248/613.full.html#related

This article cites 28 articles, 2 of which can be accessed free: /content/349/6248/613.full.html#ref-list-1

This article has been cited by 4 articles hosted by HighWire Press; see: /content/349/6248/613.full.html#related-urls This article appears in the following subject collections:

/cgi/collection/physics

Science (print ISSN 0036-8075; online ISSN 1095-9203) is published weekly, except the last week in December, by the American Association for the Advancement of Science, 1200 New York Avenue NW, Washington, DC 20005. Copyright 2015 by the American Association for the Advancement of Science; all rights reserved. The title Science is a registered trademark of AAAS.

7 AUGUST 2015 • VOL 349 ISSUE 6248 **617**